

NHS Pharmacy First – 7 clinical pathways

Urinary tract infection	Shingles	Impetigo	Infected insect bites	Acute sore throat	Acute sinusitis	Acute otitis media
Inclusion: <ul style="list-style-type: none"> Female Aged between 16 - 64 Suspected lower UTI 	Inclusion: <ul style="list-style-type: none"> 18 years and over Suspected case of shingles. Rash appeared within the last 72 hours - 7 days 	Inclusion: <ul style="list-style-type: none"> 1 year and over Signs and symptoms of impetigo Localised (4 or fewer lesions/clusters present) 	Inclusion: <ul style="list-style-type: none"> 1 year and over Infection that is present or worsening at least 48 hours after the initial bite(s) or sting(s) 	Inclusion: <ul style="list-style-type: none"> 5 years and over Suspected sore throat 	Inclusion: <ul style="list-style-type: none"> 12 years and over Suspected signs and symptoms of sinusitis Symptom duration of 10 days or more 	Inclusion: <ul style="list-style-type: none"> Aged between 1 – 17 Suspected signs and symptoms of acute otitis media
Exclusion: <ul style="list-style-type: none"> Male <16 or >64 Pregnant Breastfeeding Recurrent UTI (2 in last 6 months or 3 in last 12 months) Catheter 	Exclusion: <ul style="list-style-type: none"> < under age of 18 Pregnant or suspected pregnancy Breastfeeding with shingle sores on the breasts Shingles rash onset over 7 days ago 	Exclusion: <ul style="list-style-type: none"> < under 1 year of age Pregnancy or suspected pregnancy in individuals under 16 years of age Breastfeeding with impetigo lesion(s) present on the breast Recurrent impetigo (2 or more episodes in the same year) Widespread lesions/clusters present Systemically unwell 	Exclusion: <ul style="list-style-type: none"> < under 1 year of age Pregnancy or suspected pregnancy in individuals under 16 years of age Systemically unwell Bite or sting occurred while travelling outside the UK 	Exclusion: <ul style="list-style-type: none"> Individuals under 5 years of age Pregnancy or suspected pregnancy in individuals under 16 years of age age Recurrent sore throat/tonsillitis (7 or more significant episodes in the preceding 12 months or 5+ in each of the preceding 2 years, or 3+ in the preceding three years) Previous tonsillectomy 	Exclusion: <ul style="list-style-type: none"> Individuals under 12 years of age Pregnancy or suspected pregnancy in individuals under 16 years of age Symptom duration of less than 10 days Recurrent sinusitis ((4 or more annual episodes of sinusitis) 	Exclusion: <ul style="list-style-type: none"> Individuals under 1 year of age or over 18 years of age Pregnancy or suspected pregnancy in individuals under 16 Recurrent infection (3+ episodes in preceding 6 months, or 4+ episodes in the preceding 12 months with at least one episode in the past 6 months.)